

**POISON**

**KEEP OUT OF REACH OF CHILDREN**

**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**



**Agador<sup>®</sup>**

**Turf nematicide and miticide**

**syngenta<sup>®</sup>**

**ACTIVE CONSTITUENTS: 20 g/L ABAMECTIN**

**GROUP 6 INSECTICIDE**

For the control of nematodes and Bermudagrass Mites  
in turf as per the Directions for Use

*Imported By:*

**Syngenta Asia Pacific Pte Ltd**

No. 1 HarbourFront Avenue

#03-03 Keppel Bay Tower

Singapore 098632

Tel: +65-6333 6400

Fax: +65-6338 1594

**Emergency hotline number: +65 6334 4366**

**APVMA Approval No: 66380/102611**

**Item number, Date code**

<sup>®</sup>

## DIRECTIONS FOR USE

### Restraints

DO NOT use in the home garden  
 DO NOT apply by backpack application  
 DO NOT apply by a hand gun sprayer  
 DO NOT apply with aircraft  
 DO NOT apply through any type of irrigation system or ultra low volume spray system  
 DO NOT apply to turf under heat or moisture stress

Situation	Pests	Rate	Critical Comments
Turf	Bermudagrass Mite ( <i>Aceria cynodoniensis</i> , formerly <i>Eriophyes cynodoniensis</i> )	1 to 2 L/ha or 10 to 20 mL/100 m <sup>2</sup>	Apply AGADOR at 1 L/ha when pest pressure is low (early curative). Use the 2 L/ha rate when pest pressure is high and turf injury is already evident. Always add a non-ionic surfactant to ensure good coverage and penetration of leaf sheaths. Apply 1 to 2 consecutive AGADOR applications at 21 to 28 day intervals if population persists. <b>Refer to Application section for detailed information.</b>
	Turf-Parasitic Nematodes	2 L/ha or 20 mL/100 m <sup>2</sup>	Apply AGADOR in an early curative situation (after appropriate nematode extraction, identification and counts). Wash application in with 3 to 6 mm of irrigation or rain immediately after application. Apply up to 3 consecutive AGADOR applications at 14 to 21 day intervals. <b>Sting Nematode:</b> When counts are 100 nematodes/200 g soil or higher apply at least 3 consecutive applications of AGADOR at 14 day intervals. Switch to an alternative chemistry class if counts stay high and symptoms persist. <b>Refer to Application section for detailed information.</b>

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

**WITHHOLDING PERIOD: DO NOT GRAZE TREATED TURF/LAWN OR FEED TURF/LAWN CLIPPINGS FROM ANY TREATED AREA TO POULTRY OR LIVESTOCK**

### GENERAL INSTRUCTIONS

#### Mixing

Add the required amount of product directly to the spray tank and mix well. AGADOR mixes completely with water and should not be tank mixed with other fertilisers as it contains an iron chelate.

**Non-ionic Surfactants:** Use Agral® Spray Adjuvant (600 g ai/L) at 0.42% v/v or 420 mL/100 L spray mix. Alternatively use other non-ionic surfactants (1000 g ai/L formulations) at 0.25% v/v or 250 mL/100 L spray mix.

#### Compatibility

AGADOR is compatible with Banner® Maxx Turf Fungicide, Headway® Maxx Turf Fungicide, Heritage® Maxx Turf Fungicide, Primo® Maxx Turf Growth Regulator and Subdue® Maxx Turf Fungicide.

#### Application

Apply by ground boom sprayer or low pressure hand wand  
 DO NOT apply by backpack application  
 DO NOT apply by a hand gun sprayer  
 DO NOT apply with aircraft  
 DO NOT apply through any type of irrigation system or ultra low volume spray system  
 DO NOT apply with a nozzle height greater than 50 cm above the ground

**Nematodes:** Application volumes for root affecting nematodes should be as high as possible to ensure placement close to the soil surface. When lower application volumes are used, washing in should commence as soon as possible after application. For best results use very coarse droplets (eg Air Induction 08 nozzles) and total application volume of approximately 1000 L/ha. Preferably spray onto wet or dewy grass. Irrigate with 3 to 6 mm of water commencing within 1 hour of application.

**Bermudagrass Mite:** Application volume for Bermudagrass Mite should be adequate to ensure thorough and even coverage of the turf leaves and penetration into the crowns. Ideal application volume should be 350 to 500 L/ha. For best results use minimum coarse droplets (eg Air Induction 025 or 04 nozzles). In higher cut turf ( $\geq 15$  mm) a significant spray shielding effect can occur, impacting negatively on spray penetration and even coverage at low application volumes.

### **Insecticide Resistance Warning**

### **GROUP 6 INSECTICIDE**

For insecticide resistance management AGADOR Turf Nematicide and Miticide is a Group 6 insecticide. Some naturally occurring insect biotypes resistant to AGADOR and other Group 6 insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if AGADOR and other Group 6 insecticides are used repeatedly. The effectiveness of AGADOR on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Syngenta Australia Pty Ltd accepts no liability for any losses that may result from failure of AGADOR to control resistant insects. AGADOR may be subject to specific resistance management strategies. For further information contact your supplier, Syngenta representative or local agricultural department agronomist.

### **Integrated Pest Management**

The possible effects of AGADOR on integrated pest management (IPM) strategies in the turf industry have not been studied at the proposed rates. However, based on available information, it cannot be ruled out that AGADOR may have an adverse effect on non-target beneficial turfgrass invertebrates where such IPM is practiced.

### **PRECAUTIONS**

#### **Re-entry Period**

##### **General Public**

DO NOT allow public re-entry into treated areas until the spray has dried.

##### **Grounds Staff / Professional Operators**

**Low exposure activities:** DO NOT allow entry into treated areas until the spray has dried. When prior entry is necessary, wear chemical resistant gloves and footwear.

**High exposure activities:** DO NOT allow entry into treated areas for 4 days for high exposure activities such as hand weeding, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

### **PROTECTION OF LIVESTOCK**

DO NOT graze treated turf/lawn, or feed turf/lawn clippings from any treated area to poultry or livestock.

Highly toxic to bees. Will kill bees foraging in or near treated turf or in hives which are oversprayed or reached by spray drift. Residues may remain toxic to bees for several days after application.

### **PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

DO NOT apply under weather conditions or from spraying equipment that may cause spray to drift from the target area. Very toxic to aquatic life. DO NOT contaminate wetlands or watercourses with this product or used containers.

### **STORAGE AND DISPOSAL**

Store in the closed original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to the spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If no facility is available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with Local, State or Territory government regulations. DO NOT burn empty containers or product.

## **SAFETY DIRECTIONS**

**Harmful if swallowed. Harmful if inhaled. May irritate the eyes. Avoid contact with eyes. DO NOT inhale spray mist. When opening the container, preparing the product for use and using the product wear:**

- cotton overalls buttoned to the neck and wrist (or equivalent clothing)
- elbow length chemical resistant gloves
- half face piece respirator with organic vapour/gas cartridge or canister

**Wash hands after use. After each day's use, wash gloves, respirator (if rubber, wash with detergent and warm water) and contaminated clothing.**

## **FIRST AID**

**If poisoning occurs contact a doctor or Poisons Information Centre. Phone +65 6334 4366. If skin contact occurs, remove contaminated clothing and wash skin thoroughly.**

## **MATERIAL SAFETY DATA SHEET**

If additional hazard information is required refer to the Material Safety Data Sheet. For a copy phone +65 6333 6400

## **DISCLAIMER**

This product complies with the specifications in its statutory registration. Implied terms and warranties are excluded. Syngenta's liability for breach of the express or any non-excludable implied warranty is limited to product replacement or purchase price refund. The purchaser must determine suitability for intended purpose and take all proper precautions in the handling, storage and use of the product including those on the label and/or safety data sheet failing which Syngenta shall have no liability.

Product names marked ® or ™, the ALLIANCE FRAME  
the SYNGENTA Logo and the PURPOSE ICON  
are Trademarks of a Syngenta Group Company



\*Trademark