

# BRISKWAY

A broad-spectrum fungicide for the prevention and control of certain diseases in turf as per Direction for Use.

## Suspension Concentrate (SC)

### ACTIVE INGREDIENTS:

Azoxystrobin	18% w/w
Difenoconazole	11.3% w/w
Other ingredients:	70.7% w/w
Total:	100%

### CAUTION

**KEEP LOCKED UP AND OUT OF REACH OF CHILDREN.**

**IMPORTANT: Read the label before opening or using**

### NET CONTENT:

*Imported By:*

**Syngenta Asia Pacific Pte. Ltd.**

No. 1 HarbourFront Avenue

#03-03 Keppel Bay Tower

Singapore 098632

Tel: +65-6333 6400

Fax: +65-6338 1594

**Emergency hotline number:** +65 6334 4366

Product names marked ® or ™, the ALLIANCE FRAME  
the SYNGENTA Logo and the PURPOSE ICON  
are Trademarks of a Syngenta Group Company



## SAFETY PRECAUTIONS

Harmful if swallowed or absorbed by skin contact. Will irritate the eyes and skin. Avoid contact with skin, eyes or clothing. When opening the container and using the prepared spray, wear cotton overalls buttoned to the neck and wrist, elbow-length chemical-resistant gloves, face shield or goggles. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield/goggles and contaminated clothing. Do not contaminate water sources.

## DIRECTIONS FOR USE

Crops	Diseases Controlled	Application rate	Application interval (days)	Remarks
Zoysiagrass, warm season turfgrass	Brown patch/large patch ( <i>Rhizoctonia solani</i> )	1.5 – 2.3 l/ha	14-28	For large patch, make 1 or 2 applications prior to infections or when conditions are favourable for infection.
Seashore paspalum, warm season turfgrass	Dollar spot ( <i>Sclerotinia homoeocarpa</i> )	1.6 – 2.3 l/ha	14-21	Apply preventively when conditions are favourable for disease development. For optimum disease control, alternate Briskway with contact fungicides. If dollar spot is active, use higher rates and combine with a contact fungicide.
Bermudagrass, warm season turfgrass	Leaf spot ( <i>Bipolaris</i> spp.)	2 l/ha	14-21	Apply when conditions are favourable for disease development.

**Restricted Entry Intervals:** DO NOT allow entry into treated areas until the spray has dried unless protective clothing is worn.

## Compatibility

As formulations of other manufacturers products are beyond the control of Syngenta, all mixtures should be tested before addition to a spray tank and before application to turf to establish safety.

## FIRST AID INSTRUCTIONS

**If poisoning occurs, contact a doctor or call the Emergency Hotline number +65 6334 4366**

**If in eyes:** Rinse immediately with plenty of water, also under the eyelids, for at least 15 minutes. Remove contact lenses. Immediate medical attention is required

**If on skin:** Take off all contaminated clothing immediately. Wash off immediately with plenty of water. If skin irritation persists, call a physician.

Wash contaminated clothing before re-use.

**If inhaled:** Move the victim to fresh air.

If breathing is irregular or stopped, administer artificial respiration. Keep patient warm and at rest. Call a physician or poison control center immediately.

**If swallowed:** Seek medical advice immediately and show this container or label. Do NOT induce vomiting.

**Antidote:** There is no specific antidote available. Treat symptomatically.

## STORAGE

DO NOT contaminate water, food, or feed by storage and disposal.

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight.

## DISPOSAL

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If not recycling, triple rinse, break, crush or puncture and deliver empty packaging for appropriate disposal at an approved waste management facility. DO NOT burn empty containers or product.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

**WITHHOLDING PERIOD: DO NOT GRAZE TREATED TURF/LAWN OR FEED TURF/LAWN CLIPPINGS FROM ANY TREATED AREA TO POULTRY OR LIVESTOCK**

### **Restraints**

DO NOT apply through irrigation systems.

DO NOT spray Briskway where spray drift may reach apple trees.

DO NOT apply with spray droplet smaller than medium to coarse spray droplet size category.

DO NOT apply where wind speed is greater than 10 km/hour at application site

DO NOT apply during surface temperature inversion conditions at the application site.

DO NOT apply if there are aquatic and wetland areas including aquacultural ponds or surface streams and rivers within 5 m downwind from application areas.

### **GENERAL INSTRUCTIONS**

Briskway is a broad-spectrum product containing two fungicides. It has preventative, systemic and curative properties.

Briskway provides disease control of many major turfgrass diseases. Briskway is a Suspension concentrate (SC) formulation that may be applied as a foliar spray and integrated into a resistance management program. All applications should be made according to the use directions that follow.

### **Mixing**

Thoroughly clean spray equipment before using this product.

Make up only the amount of spray required for the immediate use.

To prepare a spray solution, partially fill the spray tank with clean water and begin agitation. Add the specified amount of Briskway to the tank, allowing time for good dispersion, then add an adjuvant, if recommended.

DO NOT use silicone based products with Briskway due to possible phytotoxicity.

Maintain agitation throughout the spraying operation. DO NOT allow spray mixture to stand overnight or for prolonged periods.

Rinse spray tank thoroughly with clean water after each day's use and dispose of pesticide rinsate by application to an already treated area.

### **Application**

Briskway may be applied with all types of spray equipment commonly used for making ground applications. Proper adjustments and calibration of spraying equipment to give good coverage is essential for good disease control.

Spray nozzles should be uniformly spaced and of the same size, and should provide accurate and uniform application. Use Syngenta's Turf XC Nozzles, or nozzles that provide a uniform droplet size and produce medium to coarse droplets (250 to 400 microns VMD).

Apply in sufficient water to provide good coverage. Typical application volumes range from 125 to 1000 L/ha. Thorough coverage is necessary to provide good disease control.

For control of foliar diseases, allow Briskway applications to completely dry prior to irrigating. For control of soil-borne diseases, Briskway can be watered in after application.

### **Efficacy**

Under certain conditions conducive to extended infection periods, use another registered fungicide for additional applications if the maximum amount of Briskway has been used. The higher rates in the rate range and/or shorter spray intervals may be required under conditions of heavy infection pressure with highly susceptible varieties, or when environmental conditions are conducive to disease.

### **Resistance Management**

Some turf disease pathogens are known to have developed resistance to products used repeatedly for their control. Briskway should be alternated with other registered fungicides that have a different mode of action and to which pathogen resistance has not developed.

**KEEP LOCKED UP OUT OF REACH OF CHILDREN AND OTHER UNAUTHORISED PERSONS, AND ANIMALS.**

**KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFF.**

### **MATERIAL SAFETY DATA SHEET**

If additional hazard information is required, refer to the Material Safety Data Sheet. For a copy phone +65 6333 6400

### **DISCLAIMER**

This product complies with the specifications in its statutory registration. Implied terms and warranties are excluded. Syngenta's liability for breach of the express or any non-excludable implied warranty is limited to

product replacement or purchase price refund. The purchaser must determine suitability for intended purpose and take all proper precautions in the handling, storage and use of the product including those on the label and/or safety data sheet failing which Syngenta shall have no liability.