

## Application Technology

### Getting Meridian to the target effectively

#### Greens – Optimal Method

Total application volume ~ 950 L/ha

Turbo FloodJet TF5 or TurfJet TTJ10

5 km/h @ 3 bar pressure

**Wash in with at least 6 mm of irrigation**

#### Fairways – Optimal Method

Total application volume ~ 590 L/ha

Turbo FloodJet TF5 or TurfJet TTJ10

8 km/h @ 3 bar pressure

**Wash in with at least 6 mm of irrigation**



Save up to 60,000 litres of water per hectare

Irrigation savings

Greater availability in dry conditions

Season long control of grubs



**syngenta**

[www.syngentaturf.sg](http://www.syngentaturf.sg)

For further information contact Syngenta  
Technical Product Advice Line at +65  
6333 6400

Product names marked ® and the Syngenta logo are trademarks of a Syngenta Group Company.  
\*Registered trademark. ABN 33 002 933 717 Syngenta Crop Protection Pty Limited 2-4 Lyonpark Road, North Ryde, NSW 2113. The information contained in this brochure is believed to be accurate. No responsibility is accepted in respect of this information, save those non-excludable conditions implied by any Federal or State legislation or law of Territory. SYN07/226.



Meridian


Imidacloprid

8FD1315\_revised\_L0707




## Irrigation savings “washing in” after application

Meridian® can help to save up to 60,000 litres of water per hectare during post grub-insecticide application “wash in”. The superior solubility of the active ingredient – thiamethoxam (neonicotinoid) – is the main reason for this benefit. Thiamethoxam is approximately 8 times more soluble than imidacloprid (Fig. 1), thus increasing robustness and water savings during incorporation.



## Effective in wet and dry soil conditions

Neonicotinoids age in soil, with the majority of the active ingredient available during the first 30 days dissolved in soil moisture. Increased solubility of thiamethoxam during this time will ensure elevated levels of availability and thus efficacy in drier soil conditions. In the longer term (20-100 days), thiamethoxam will bind to soil, to be released more readily back (desorbed) into solution than imidacloprid. This is called bio-availability, meaning the active ingredient is available to the biological system – plants for uptake and/or contact with burrowing grubs. Higher bio-availability contributes to higher levels of efficacy. Meridian will ensure high level performance in most soil conditions (Fig. 2). Meridian may therefore perform better than imidacloprid in drier soils such as fairways, semi-roughs, ovals and parks lacking regular irrigation (Fig. 3)



## Figure 3 Performance in dry soil

	0-30 days	20-100 days
	Dissolved (majority of active)	Soil bound (majority of active)
<b>Meridian</b>	Higher solubility Increased bio-availability and efficacy	Easier desorption Increased bio-availability and efficacy
<b>imidacloprid</b>	Lower solubility Decreased bio-availability and efficacy	Difficult desorption Decreased bio-availability and efficacy

## Season-long control of grubs

### Scarabs (African Black Beetle)

The African Black Beetle (*Heteronychus arator*) (Fig. 4) occurs in all states and completes a single life cycle per year. Mating occurs in early spring with egg laying activities peaking during October in the northern states (QLD, NSW and WA) and a little later during early November in the southern states (ACT, VIC, TAS and SA) (Fig. 6). First and second instar grubs feed on roots in the upper soil layers. The excellent contact and systemic action of Meridian controls grubs and protects roots.



Figure 4

### Turfgrass Weevils (Billbug or La Plata Weevil)

The Billbug Weevil (*Sphenophorus brunnipennis*) (Fig. 5) occurs in all states and completes multiple lifecycles – commonly 2 and exceptionally 3 per year. The first generation overlaps largely with the scarabs during spring and early summer. The second generation normally peaks with egg laying during mid summer (January)(Fig. 6). This generation is normally at lower infestation levels, but still has a high enough impact to warrant treatment. On the rare occasion that a third generation occurs, it peaks with egg laying during early autumn (March-April) (Fig. 6). First instar larvae initially feeds in the stems, before dropping to the soil, continuing feeding on roots in the upper soil layers. The excellent systemic action of Meridian within plants ensure early control of this pest, even prior to dropping to the soil.



Figure 5

Meridian has excellent efficacy on African Black Beetle and Billbug larvae (1st and 2nd instar only). The best timing for application is thus during or shortly after peak egg laying. The illustration below suggests the optimum time of application to ensure optimal results. Meridian has varied dose rates to accommodate the need for a follow up application when the second and/or third generations of Billbug is to be controlled.

## Figure 6 Season long grub control program

